



Monoclonal antibody to C-reactive protein, clone 5F5, hIgG1


Catalogue #	R1-259-100
Immunogen:	Recombinant C-reactive protein
Immunogen Description:	Recombinant C-reactive protein
Source:	Human
Clonality:	Human monoclonal
Clone:	5F5
Class:	hIgG1
Application:	ELISA, CLIA
Kd:	2.249 x 10 ⁻¹⁰ M
Purification:	Produced recombinantly using CHO-based cell line (expressed by QMCF technology). Purified using protein A affinity chromatography followed by desalting
Purity:	>95%
Concentration:	1 mg/ml
Buffer:	PBS, pH 7.4
QC:	LabChip protein analysis, analytical SEC, Octet binding analysis
Shipping:	Shipped with blue ice.
Storage:	Store at +4 °C. Avoid multiple freeze-thaw cycles.



Figure 1. CE-SDS virtual gel output (LabChip GX) for monoclonal antibody to C-reactive protein, clone 5F5 under non-reduced (NR, left) and reduced (R, right) conditions.

Peak #	RT (min)	Estimated Mw (Da)*	Area	Area %
1	7.615	340464	22.10	0.83
2	8.727	150864	2626.31	99.17

Chromatogram




*Calculated using calibration curve obtained from AdvanceBio SEC 300A Protein Standard (p/n 5190-9417) retention times. Peaks with integrated areas below 0.5% of the calculated values were excluded from analysis.

Figure 2. Analytical SEC of final product.

Peak #	RT (min)	Estimated Mw (Da)*	Area	Area %
1	7.602	342617	26.54	0.96
2	8.727	150846	2738.14	99.04

Chromatogram



*Calculated using calibration curve obtained from AdvanceBio SEC 300A Protein Standard (p/n 5190-9417) retention times. Peaks with integrated areas below 0.5% of the calculated values were excluded from analysis.

Figure 3. HPLC analytical SEC after 3 freeze-thaw cycles.




Figure 4. Octet binding analysis, antibody was loaded on sensor for capture of C-reactive protein in different concentrations.