



## Monoclonal antibody to C-reactive protein, clone 6F6, hIgG1


Catalogue #	R1-260-100
Immunogen:	Native human C-reactive protein
Immunogen Description:	Native human C-reactive protein purified from human pleural fluid.
Source:	Human
Clonality:	Human monoclonal
Clone:	6F6
Class:	hIgG1
Application:	ELISA, CLIA
Kd:	$3.570 \times 10^{-11}$ M
Purification:	Produced recombinantly using CHO-based cell line (expressed by QMCF technology). Purified using protein A affinity chromatography followed by desalting
Purity:	>90%
Concentration:	1 mg/ml
Buffer:	PBS, pH 7.4
QC:	LabChip protein analysis, analytical SEC, Octet BLI analysis
Shipping:	Shipped with dry ice.
Storage:	Store at +4 °C.



**Figure 1.** CE-SDS virtual gel output (LabChip GX) for monoclonal antibody to C-reactive protein, clone 6F6 under non-reduced (NR, left) and reduced (R, right) conditions.

Peak #	RT (min)	Estimated Mw (Da)*	Area	Area %
1	7.420	359333	28.11	1.03
2	8.908	131568	2699.35	98.97

## Chromatogram




\*Calculated using calibration curve obtained from AdvanceBio SEC 300A Protein Standard (p/n 5190-9417) retention times. Peaks with integrated areas below 0.5% of the calculated values were excluded from analysis.

**Figure 2.** Analytical SEC of final product.


Peak #	RT (min)	Estimated Mw (Da)*	Area	Area %
1	7.434	350364	21.11	1.03
2	8.907	132648	2020.68	98.97

## Chromatogram



\*Calculated using calibration curve obtained from AdvanceBio SEC 300A Protein Standard (p/n 5190-9417) retention times. Peaks with integrated areas below 0.5% of the calculated values were excluded from analysis.

**Figure 3.** HPLC analytical SEC after 3 freeze-thaw cycles.



**Figure 4.** Octet BLI analysis, antibody was loaded on sensor for capture of C-reactive protein in different concentrations.