



Mouse mAb to HCV subtype 1b NS5B (clone 10D6)

Catalogue #	A2-425-100
Immunogen:	HCV 1b NS5B
Immunogen Description:	Recombinant Hepatitis C Virus subtype 1b non-structural protein 5B (NS5B) RNA-dependent RNA polymerase (RdRp)
Uniprot ID:	Q9WMX2
Clonality:	Mouse monoclonal
Clone:	10D6
Class:	mIgG1
Reactivity:	Human, HCV subtype 1b NS5B, Epitope mapped to amino acids 111-130 (LSSKAVNHIRSVWKDLLEDT)
Application:	ELISA, WB, IP
Protocol:	Optimal conditions for IP should be determined for each particular application.
ELISA:	0,25 µg/ml
IF:	-
Purification:	Protein G purification
Buffer:	PBS pH 7.4, with 0.1% sodium azide
Related Products:	HCV subtype 1b monoclonal antibodies. For more information visit www.icosagen.com/products/?antibodies
Shipping:	This product is shipped in non-frozen liquid form in ambient conditions
Storage:	Store at -20 ...-70 °C upon receipt. Divide antibody into aliquots prior usage. Avoid multiple freeze-thaw cycles as product degradation may result.
Background:	Non-structural protein 5B (NS5B) represents the RNA-dependent RNA polymerase (RdRp) of Hepatitis C Virus, which is a small positive strand

RNA virus in the family Flaviviridae. HCV is a major causative agent of acute and chronic hepatitis, hepatocellular carcinoma and liver cirrhosis. The single subunit RNA-dependent RNA polymerase is absolutely essential for the viral replication.

References

Nikonov A, Juronen E and Ustav M (2008). Functional characterization of fingers subdomain-specific monoclonal antibodies inhibiting the hepatitis C virus RNA-dependent RNA polymerase. J. Biol Chem. 283(35):24089-102.




Figure 1. IP was carried out with NS5B specific mAb 10D6 using the lysates of Huh7 cells harboring selectable subgenomic HCV RNA replicon (upper panel) or plain Huh7 cells (lower panel). NS5B polyclonal antibodies (a-NS5B) and a-Bcl2 mAb, directed against cellular protein, were used as positive and negative controls respectively. Asterisk indicates immunoglobulin heavy chain; protein blots were probed with NS5B specific mAb 10D6




Figure 2. WB analysis of Huh-7 cell line expressing NS5B protein. Detected by 10D6 antibody. Line 1. Huh-7 transfected with HCV replicon, Line 2. Huh-7 non-transfected cells.