


## Mouse mAb to hNRTN (clone 4G8)

| Catalogue #               | 321-100  |
|---------------------------|--|
| Immunogen:                | human Neurturin  |
| Immunogen<br>Description: | Recombinant human NRTN protein produced<br>using CHO-based Icosagen Cell factory Ltd.<br>proprietary suspension cell line. Protein was<br>purified from the cell culture supernatant |
| Alternative Names:        | NRTN, NTN  |
| Uniprot ID:               | Q99748   |
| Clonality:                | Mouse monoclonal   |
| Clone:                    | 4G8  |
| Class:                    | mlgG1  |
| Reactivity:               | Human NRTN   |
| Application:              | ELISA, WB, IF  |
| Protocol:                 | Monoclonal antibody working titer has to be established practically for each particular antigen and assay format   |
| ELISA:                    | 0,05-0,1 μg/ml   |
| IF:                       | 2-10 μg/ml   |
| Purification:             | Protein G purification   |
| Buffer:                   | PBS pH 7.4, with 0.1% sodium azide   |
| Shipping:                 | This product is shipped in non-frozen liquid form in ambient conditions  |
| Storage:                  | Store at – 20 or -70 °C upon receipt. Divide<br>antibody into aliquots prior usage. Avoid multiple<br>freeze-thaw cycles as product degradation may<br>result                        |
| Background:               | Neurturin is a member of the GDNF family of ligands. GDNF family proteins are distant members of the TGF-beta superfamily  |



**Figure 1. Western-Blot detection of human NRTN expressed in CHO cells.** Line 1. hNRTN-containing CHOEBNALT85 cell culture supernatant. Line 2. Negative control – CHOEBNALT85 cell culture supernatant. 5 µl of supernatant were loaded per line. Primary antibody dilution 1:5000 was used. AP-conjugated Goat-anti mouse, was used as secondary antibody. Antibody recognizes protein dimeric, and monomeric variants