


## Mouse mAb hGDNF (clone 3C1)


Catalogue #	315-100
Immunogen:	Human GDNF
Immunogen Description:	Recombinant human GDNF protein produced using <i>E. coli</i> expression system
Alternative Names:	Astrocyte-derived trophic factor (ATF)
Uniprot ID:	P39905
Clonality:	Mouse monoclonal
Clone:	3C1
Class:	mIgG1
Reactivity:	Human GDNF
Application:	ELISA, WB, IF, IHC
Protocol:	Monoclonal antibody working amount has to be established practically for each particular antigen and assay format
ELISA:	50-100 ng/ml
IF:	2-10 µg/ml
IHC:	5 µg/ml
Purification:	Protein G purification
Buffer:	PBS pH 7.4, with 0.1% sodium azide
Shipping:	This product is shipped in non-frozen liquid form in ambient conditions
Storage:	Store at -20...-70°C upon receipt. Divide antibody into aliquots prior usage. Avoid multiple freeze-thaw cycles
Background:	GDNF is a neurotrophic factor that enhances survival and morphological differentiation of dopaminergic neurons and increases their high-affinity dopamine uptake. Ligand for the GFR-alpha-3-RET receptor complex but can also

activate the GFR-alpha-1-RET receptor complex




**Figure 1. Western Blot testing of GDNF monoclonal antibody 3C1.** Analysis was performed with antiGDNF monoclonal antibody 3C1 HRP conjugate. 15 µl of CHOEBNALT85 GDNF producing cell culture supernatant was loaded per lane. Analysis was performed in reducing (A) and non-reducing (B) conditions

A




**B**




**Figure 2. Immunofluorescence detection of human GDNF expressed in U2OS cells by anti-GDNF monoclonal antibody 3C1.** Anti-GDNF antibody 3C1 concentration in IF experiment was 10  $\mu\text{g/ml}$ . Goat anti-mouse AlexaFluor488 was used as secondary antibody. For nuclear staining DAPI was used. ArrayScan VTI platform (Thermo Scientific) was used for image acquisition (10x objective). Composite picture was generated using pseudocolors green for GDNF specific signal and blue for nuclei. A. GDNF-expressing U2OS cells; B. Negative control (non-transfected U2OS cells)

**A**



**B**



**Figure 3. Immunohistochemistry testing of anti-GDNF monoclonal antibody**

**3C1.** Analysis was performed using formalin-fixed paraffin-embedded human cerebral cortex tissue sections from Alzheimer's disease patients. Tissue sections were boiled with sodium citrate buffer (pH 6) for antigen retrieval. Incubation with primary antibody at 5 µg/ml was performed overnight at 4°C. DAKO EnVision™ Detection System, Peroxidase/DAB was used for visualization. Sections were counterstained with toluidine blue and mounted with Eukitt mounting medium. A. GDNF staining by monoclonal antibody 3C1; B. Negative staining without primary antibody