


## Human IgG1-kappa antibody to SARS CoV-2 S1 RBD (clone 38G5)

Catalogue #	R1-159-100
Immunogen:	SARS-CoV-2 virus
Immunogen Description:	SARS-CoV-2 virus (PBMCs isolated from patients recovered from SARS-CoV-2 infection, antibodies discovered by using HybriFree Technology).
Clonality:	Human monoclonal
Clone:	38G5
Class:	hIgG1
Reactivity:	SARS-CoV-2 Spike RBD
Dissociation constant ( $K_D$ ):	$< 1.0 \times 10^{-12}$ M (measured against SARS-CoV-2 Trimeric Spike protein) $3.46 \times 10^{-8}$ M (measured against SARS-CoV-2 Spike S1 protein)
Application:	ELISA
ELISA:	0,02-0,04 ng/ml
Purification:	Protein A affinity chromatography following desalting
Buffer:	PBS pH 7.4
Shipping:	Shipped at ambient temperature.
Storage:	Store at +4 °C. Avoid multiple freeze-thaw cycles.




**Figure 1.** Coomassie-stained SDS-PAGE analysis of monoclonal antibody to SARS-CoV-2 Spike RBD, clone 38G5. 4-12% gradient gel is used for analysis. Lane 1. Monoclonal antibody to SARS-CoV-2 Spike RBD, clone 38G5 (-DTT). Lane 2. Size Marker. Lane 3. Monoclonal antibody to SARS-CoV-2 Spike RBD, clone 38G5 (+DTT). Lane 4. Size Marker.

## Peak Table

Peak #	RT (min)	Area	Area %
1	7.494	2356.85	100.00

## Chromatogram




**Figure 2.** HPLC analytical SEC for final product.


## Peak Table

Peak #	RT (min)	Area	Area %
1	7.490	1605.93	100.00

## Chromatogram



**Figure 3.** HPLC analytical SEC after 3 freeze-thaw cycles.



**Figure 4.** Octet RED96e analysis, antibody was loaded on sensor for capture of Spike S1 protein.